



## Cogmed Working Memory Training

### Research Evidence for the Improvement of Working Memory and Attention

#### Summary

**Working memory (WM) is a key cognitive ability, necessary for control of attention and academic performance. Cogmed Working Memory Training (CWMT) is a researched-based, systematic method for improving WM through computerized training. The improvements after CWMT transfers to non-trained WM tasks, an effect equivalent of about 2 years of normal childhood development. The effect on attention is documented in healthy adults as well as in patient groups, including long-term follow-up studies. In particular, the clinical research has documented improvement of the key inattentive symptoms in:**

- **Children with ADHD**
- **Children who have received cancer treatment**
- **Adults after stroke**

**Improvements in mathematics has been documented in several trials but is larger in typically developing children than in children with WM impairments. There are now more than 120 studies of CWMT, by independent research groups around the world, which makes it the most well-researched method of improving WM and attention.**

#### What is Working Memory?

Working Memory (WM) is our ability to keep information online. We use WM to keep relevant information and plans for what to do next in mind when solving problems, or completing tasks. WM is closely related to control of attention, and these abilities make use of partly the same regions in the brain: the parietal and prefrontal cortices (Ikkai and Curtis, 2010; Constantinidis and Klingberg, 2016), areas that are targeted by CWMT (Olesen et al., 2004). One way to put it is that we use WM to remember what to focus on.

All individuals vary in their WM capacity. A low WM can be hereditary, linked to environmental effects such as stress, or caused by injury to the brain. Many diagnoses and clinical conditions are associated with low WM and attention deficits, including ADHD, ADD, traumatic brain injury, premature birth, stroke and cancer treatment.

The results of an impaired WM are similar, independent of the cause: an inability to remember plans and instructions, and to pay attention to the task at hand, which is often perceived as the individual being distracted. This is especially pronounced during mentally demanding situations. In children, WM deficits are associated with academic problems in both reading and mathematics.

### Cogmed Improves Working Memory Capacity and Attention

Cogmed is based on research showing that WM capacity can be increased through training. This was originally shown in studies from the Karolinska Institute (Klingberg et al., 2002; Klingberg et al., 2005), and has since been replicated by independent research groups around the world. There are now more than 120 published studies of CWMT.

The key finding is that WM capacity is improved not only for the tasks that are part of the training program, but also for tasks that are dissimilar. For example, studies at Cambridge University, UK and Karolinska Institute, Sweden, show that CWMT improves the ability of children to remember and perform long and WM-demanding verbal instructions, a task that is highly relevant in daily life (Holmes et al., 2009a; Bergman-Nutley and Klingberg, 2014). Another example is transfer to a test battery (Automated WM Assessment) developed to specifically measure complex WM (Holmes et al., 2009b; Carlson-Green et al., 2017; Peers et al., 2020). A third example is improved ability to keep in mind, update and add digits (The Paced Auditory Serial Addition Task), a task not part of the training (Lundqvist et al., 2010; Brehmer et al., 2012). Altogether, this means that CWMT **transfers to non-trained WM tasks.**

A summary of the improvements in 12 CWMT studies found significant, long-lasting improvements in WM task performance: 0.7 standard deviations compared to the control groups (Spencer-Smith and Klingberg, 2015). This is larger than the effect of physical exercise on cognition (Verburgh et al., 2014), or of stimulant medication on WM capacity (Roberts et al., 2020).

Another way to illustrate the impact on cognition is to compare it to years of development. When children grow older, their WM improves by around 0.25 SD per year for a 10-year-old child (Ullman et al., 2014). The 0.7 SD deviation **improvement from CWMT is comparable to at least 2 years of normal development.**

Because WM and attention are closely related (Lui and Tannock, 2007; Ikkai and Curtis, 2010), we would expect that improvement of WM also leads to improvements in attention during neuropsychological tasks as well as in everyday life.

In a meta-analysis of 12 CWMT studies, improvement in attentive behavior in everyday life was documented independent of diagnosis and age, an effect that was retained at

follow-up (Spencer-Smith and Klingberg, 2015). This includes a study in 100 healthy young and older adults with significant improvements documented with both neuropsychological tests and questions about attention problems in everyday life (Brehmer et al., 2012).

The mean effect size of this improvement ( $d = 0.4$ ) is stronger than, for example, the effects of SSRI on depression (Kirsch et al., 2008).

### Cogmed WM Training Improves Attention in Children with ADHD

Clinical studies has mainly been focused on ADHD. Table 1 summarizes six published studies (Klingberg et al., 2005; Gropper and Tannock, 2009; Beck et al., 2010; Green et al., 2012; Egeland et al., 2013; Bigorra et al., 2015), documenting how CWMT improves attention in ADHD, compared to control groups. Five studies are from children, one from adults. The questions used for estimating attention are similar or identical to those used to diagnose inattention problems, in school and at home, in ADHD. This means that **CWMT transfers to better attention in everyday life in children with ADHD**. One of the largest studies, by Bigorra and collaborators, from the University Hospital in Barcelona, concluded:

*“CWMT had a significant impact on ADHD deficits by achieving long-term far-transfer effects.”*

Study	N	Age	Ctrl group	Inattention sx
Klingberg, et al. (2005)	53	7-12	Active	DSM-IV $d = 0.7$
Beck, et al. (2010)	52	7-17	Wait-list	DSM-IV $d = 0.76$
Green et al. (2012)	26	7-14	Active	RAST off-task $d = 1.2$
Egeland 2013	67	10-12	Wait-list	ARS attention $d = 0.5$ (par) 0.4 (teach) ns
Gropper et al. (2014)	62	19-52	Wait-list	ASRS $d = 0.42$
Bigorra et al. (2015)	66	7-12	Active	ADHD composite score $d = 0.39$ (par) 0.69 (teach)

Table 1. Controlled studies showing improved attention in children with ADHD. DSM-IV = diagnostic questions of inattention symptoms from the Diagnostic and Statistical Manual, version 4; RAST=Restricted Academic Setting Task; ARS=ADHD Rating Scale; ASRS=Adult ADHD Self-Report Scale.

There has been one larger study that did not find a similar effect (Chacko et al., 2014). This study included children from low socioeconomic status regions, where a majority had Oppositional Defiant Disorder (ODD). Therefore, we do not recommend CWMT if ODD is part of the problem. CWMT is more likely to succeed in individuals where the deficit in WM and attention is the main problem.

## Cogmed Enhances Working Memory and Attention in Children Treated for Cancer

Children who have been treated with cytostatic and/or radiation therapy towards the brain suffer cognitive problems due to the effect of the treatment on the growing brain. Impaired working memory is a key component, associated with inattention and lower academic performance.

A series of studies from George Washington University School of Medicine, Washington, and St Jude Children’s Research Hospital, Memphis, has evaluated CWMT in controlled studies and found significant, long-lasting improvements in WM and attention, including cognitive symptoms in everyday life (Hardy et al., 2013; Conklin et al., 2015; Carlson-Green et al., 2017; Conklin et al., 2017) (Table 2). In addition, they noted significant improvements in learning. Dr. Heather Conklin et al (2015) concluded their research paper:

*“Study findings show computerized cognitive training is feasible and efficacious for childhood cancer survivors, with evidence for training-related neuroplasticity.”*

Study	N	Age	Ctrl group	Inattention sx	Other behavior	Cognition
Hardy 2012	20	8-16	Active Randomized	CPRS-3 inatt. d = 0.4	CPRS-3 Learning d = 0.8	WRAML2 d = 0.9
Conklin 2015	68	10-14	Waitlist Randomized	CPRS inatt d = 0.7	CPRS-3 Exec funct d = 0.8	WISC d = 0.8
Conklin 2017	62	10-14	Waitlist Randomized	CPRS. Effects maintained at 6 m	Effects maintained at 6 m.	Effects maintained at 6 m.
Carlsson-Green 2017	20	8-18	No ctrl group	CBCL 6 m. after intervention	BRIEF p<0.01 6 m. after intervention	AWMA WJ-III math

Table 2. Studies of pediatric cancer. CPRS = Conner’s Parents Rating Scale; WRAML2=Wide Range Assessment of Memory and Learning; WISC=Wechsler Intelligence Scale for Children; CBCL= Achenbach Child Behavior Check List; BRIEF=Behavior Rating Inventory of Executive Function; AWMA=Automated Working Memory Assessment; Woodcock-Johnson Tests of Achievement

## Cogmed Enhances Working Memory and Attention in Stroke Patients

Stroke often results in cognitive problems, where impaired WM is a key component. A series of studies has documented how CWMT improves cognitive performance in stroke patients, including up to 6 months' follow-up measures (Westerberg et al., 2007; Lundqvist et al., 2010; Johansson and Tornmalm, 2012; Akerlund et al., 2013; Bjorkdahl et al., 2013; Peers et al., 2020). Åkerlund and colleagues, from the Department of Rehabilitation Medicine, Sahlgrenska University Hospital, Gothenburg, Sweden, concluded:

*“Results indicated that computerized WM training can improve working memory, cognition and psychological health.”*

Study	N	Age	Ctrl group	Inattention	Other	Followup
Westerberg,et al (2007)	18	34-65	Wait-list, rnd	CFQ	PASAT, Ruff, WM	
Lundqvist et al. (2010)	21	20-65	Wait-list, rnd, xo	Occupational perf	PASAT, list span	4,20 w
Johansson et al. (2012)	18	17-64	No ctrl gr	CFQ, occup perf		6 month
Åkerlund et al. (2013)	47	22-63	Wait-list, rnd, xo	Psychological health	BNIS cog test	6, 18 w
Björkdahl et al. (2013)	47	22-63	Wait-list, rnd, xo	Cognitive question	Digit-span	6, 18 w
Peers et al. (2018)	23	28-74	Wait-list, rnd, xo	Core symptoms	AWMA, TVA	No follow

Table 3. Studies of adult stroke patients. CFQ = Cognitive Failure Questionnaire; PASAT = Paced Auditory Serial Addition Task; Ruff=The Ruff Selective Attention Test; BNIS=Barrow Neurological Institute Screen for Higher Cerebral Functions; AWMA = Automated WM Assessment; TVA=Theory of Visual Attention; rnd = randomized; xo = cross-over.

### Potential Relevance of CWMT for Academic Performance in Typically Developing Children

WM capacity is highly correlated with performance in mathematics and reading. But, in contrast to attention, these abilities are also dependent on many other factors, including knowledge stored in long-term memory. It is therefore likely that it is more difficult to improve math than attention through WM training.

There are several studies documenting improvements in mathematical performance after CWMT, the two largest being a study of 155 children with self-perceived WM and attention problems and a control-group of 304 children (Bergman-Nutley and Klingberg, 2014). Here, repeated tests of arithmetic in both groups showed a gradual improvement during training compared to the control group.

A study of 572 typically developing children, randomized to training or teaching as usual, researchers from University of Mainz and Zürich (Berger, 2020) found that improvement in mathematical performance directly after training was modest (effect size 0.2), but when children were followed up with a test one year after intervention, the difference between intervention and control group had increased to 0.4, which is equivalent to about half a year of normal improvement in primary school and 1 year of improvement in secondary school. Berger et al. note that the improvement in mathematics after training is about twice as large as the improvement seen after reducing the number of students in each classroom. This finding is in line with a smaller study from the universities of Exeter and Cambridge, which found that Cogmed training in 23 typically developing children significantly improved mathematical reasoning .

There is one large study which failed to find improvements in mathematics (Roberts et al., 2016) which included children who were selected based on their low-WM (15<sup>th</sup> percentile). At this stage, a plausible hypothesis is that improvements in mathematics after CWMT is smaller than that of the effect on inattention, develops gradually and depends on the child's WM capacity, where the typical population gain more than children with low WM. Children with low WM will likely need about twice as much training to have the same gains in mathematics as children with a typical WM capacity.

Berger and colleagues concluded their paper:

*“We find not only substantial near transfer effects on WM capacity that emerge right after the five-week training period and last throughout all evaluation waves; we also find far transfer effects on several important skills—geometry, reading ability, a measure of fluid IQ, children’s ability to inhibit pre-potent impulses, and teacher-rated self-regulation ability”*

*“Moreover, these far-transfer effects emerge over time and only become fully visible after 12-13 months. Finally, we document that 3–4 years after the intervention, the children who received training have a roughly 16 percentage points higher probability of entering the academic track in secondary school.”*

### Methodological notes on effect sizes and power

In a meta-analysis of CWMT on ratings of attention an average effect size of 0.4 was found (Spencer-Smith and Klingberg, 2015). Assuming that this is representative a study needs to include at least 200 subjects (100 training and 100 control) to have adequate statistical power (>80% chance of finding an effect). Studies with 60 subjects (30 in each group) has a statistical power of around 0.35, i.e. we would expect that around 6-7 out of 10 studies do not find a p-value less than 0.05 even if the training effect is real. To have adequate power to detect an effect size of 0.3, around 350 subjects (175 in each group) is needed. Looking back at the last 20 years of studies about cognitive interventions, we can observe that most studies have been underpowered.

Research on the benefits of cognitive training, including CWMT, will continue for several years. But CWMT is at this point the most well researched training method in for improving WM and attention.

## Key references

- Akerlund E, Esbjornsson E, Sunnerhagen KS, Bjorkdahl A (2013) Can computerized working memory training improve impaired working memory, cognition and psychological health? *Brain Inj* 27:1649-1657.
- Beck SJ, Hanson CA, Puffenberger SS, Benninger KL, Benninger WB (2010) A controlled trial of working memory training for children and adolescents with ADHD. *J Clin Child Adolesc Psychol* 39:825-836.
- Berger EMF, E.; Hermes, H.; Schunk, D.; Winkel, K.; (2020) The impact of working memory training on children's cognitive and noncognitive skills. In: SSRN.
- Bergman-Nutley S, Klingberg T (2014) Effect of working memory training on working memory, arithmetic and following instructions. *Psychol Res* 78:869-877.
- Bigorra A, Garolera M, Guijarro S, Hervas A (2015) Long-term far-transfer effects of working memory training in children with ADHD: a randomized controlled trial. *Eur Child Adolesc Psychiatry*.
- Bjorkdahl A, Akerlund E, Svensson S, Esbjornsson E (2013) A randomized study of computerized working memory training and effects on functioning in everyday life for patients with brain injury. *Brain Inj* 27:1658-1665.
- Brehmer Y, Westerberg H, Backman L (2012) Working-memory training in younger and older adults: training gains, transfer, and maintenance. *Front Hum Neurosci* 6:63.
- Carlson-Green B, Puig J, Bendel A (2017) Feasibility and efficacy of an extended trial of home-based working memory training for pediatric brain tumor survivors: a pilot study. *Neurooncol Pract* 4:111-120.
- Chacko A, Bedard AC, Marks DJ, Feirsen N, Uderman JZ, Chimiklis A, Rajwan E, Cornwell M, Anderson L, Zwilling A, Ramon M (2014) A randomized clinical trial of Cogmed Working Memory Training in school-age children with ADHD: a replication in a diverse sample using a control condition. *J Child Psychol Psyc* 55:247-255.
- Conklin HM, Ashford JM, Clark KN, Martin-Elbahesh K, Hardy KK, Merchant TE, Ogg RJ, Jeha S, Huang L, Zhang H (2017) Long-Term Efficacy of Computerized Cognitive Training Among Survivors of Childhood Cancer: A Single-Blind Randomized Controlled Trial. *J Pediatr Psychol* 42:220-231.
- Conklin HM, Ogg RJ, Ashford JM, Scoggins MA, Zou P, Clark KN, Martin-Elbahesh K, Hardy KK, Merchant TE, Jeha S, Huang L, Zhang H (2015) Computerized Cognitive Training for Amelioration of Cognitive Late Effects Among Childhood Cancer Survivors: A Randomized Controlled Trial. *J Clin Oncol* 33:3894-3902.
- Constantinidis C, Klingberg T (2016) The neuroscience of working memory capacity and training. *Nature Reviews Neuroscience*.
- Egeland J, Aarli AK, Saunes BK (2013) Few Effects of Far Transfer of Working Memory Training in ADHD: A Randomized Controlled Trial. *PLoS One* 8.
- Green CT, Long DL, Green D, Iosif AM, Dixon JF, Miller MR, Fassbender C, Schweitzer JB (2012) Will working memory training generalize to improve off-task behavior in children with attention-deficit/hyperactivity disorder? *Neurotherapeutics* 9:639-648.
- Gropper RJ, Tannock R (2009) A pilot study of working memory and academic achievement in college students with ADHD. *Journal of attention disorders* 12:574-581.
- Hardy KK, Willard VW, Allen TM, Bonner MJ (2013) Working memory training in survivors of pediatric cancer: a randomized pilot study. *Psychooncology* 22:1856-1865.
- Holmes J, Gathercole SE, Dunning DL (2009a) Adaptive training leads to sustained enhancement of poor working memory in children. *DevSci* 12:F9-15.
- Holmes J, Gathercole SE, Place M, Hilton KA, Dunning DL, Klingberg T (2009b) Working memory deficits can be overcome: impacts of training and medication on working memory in children with ADHD. *Appl Cogn Psychol* 12:9-15.
- Ikkai A, Curtis CE (2010) Common neural mechanisms supporting spatial working memory, attention and motor intention. *Neuropsychologia*.
- Johansson B, Tornmalm M (2012) Working memory training for patients with acquired brain injury: effects in daily life. *Scand J Occup Ther* 19:176-183.

- Jones, S.J., Milton, F, Mostazir, M, Adlman, A.R. (2020) The academic outcomes of working memory and metacognitive strategy training in children: a double-blind randomized controlled trial. *Dev. Sci.* DOI:10.1111/desc.12870.
- Kirsch I, Deacon BJ, Huedo-Medina TB, Scoboria A, Moore TJ, Johnson BT (2008) Initial severity and antidepressant benefits: a meta-analysis of data submitted to the Food and Drug Administration. *PLoS Med* 5:e45.
- Klingberg T, Forssberg H, Westerberg H (2002) Training of working memory in children with ADHD. *J Clin Exp Neuropsychol* 24:781-791.
- Klingberg T, Fernell E, Olesen PJ, Johnson M, Gustafsson P, Dahlstrom K, Gillberg CG, Forssberg H, Westerberg H (2005) Computerized training of working memory in children with ADHD--a randomized, controlled trial. *J Am Acad Child Adolesc Psychiatry* 44:177-186.
- Lui M, Tannock R (2007) Working memory and inattentive behaviour in a community sample of children. *Behav Brain Funct* 3:12.
- Lundqvist A, Grundstrom K, Samuelsson K, Ronnberg J (2010) Computerized training of working memory in a group of patients suffering from acquired brain injury. *Brain Inj* 24:1173-1183.
- Olesen PJ, Westerberg H, Klingberg T (2004) Increased prefrontal and parietal activity after training of working memory. *Nat Neurosci* 7:75-79.
- Peers PV, Astle DE, Duncan J, Murphy FC, Hampshire A, Das T, Manly T (2020) Dissociable effects of attention vs working memory training on cognitive performance and everyday functioning following fronto-parietal strokes. *Neuropsychol Rehabil* 30:1092-1114.
- Roberts CA, Jones A, Sumnall H, Gage SH, Montgomery C (2020) How effective are pharmaceuticals for cognitive enhancement in healthy adults? A series of meta-analyses of cognitive performance during acute administration of modafinil, methylphenidate and D-amphetamine. *Eur Neuropsychopharmacol* 38:40-62.
- Roberts G, Quach J, Spencer-Smith M, Anderson PJ, Gathercole S, Gold L, Sia KL, Mensah F, Rickards F, Ainley J, Wake M (2016) Academic Outcomes 2 Years After Working Memory Training for Children With Low Working Memory: A Randomized Clinical Trial. *JAMA Pediatr* 170:e154568.
- Spencer-Smith M, Klingberg T (2015) Benefits of a working memory training program for inattention in daily life: a systematic review and meta-analysis. *PLoS One* 10:e0119522.
- Ullman H, Almeida R, Klingberg T (2014) Structural maturation and brain activity predict future working memory capacity during childhood development. *J Neurosci* 34:1592-1598.
- Verburgh L, Konigs M, Scherder EJA, Oosterlaan J (2014) Physical exercise and executive functions in preadolescent children, adolescents and young adults: a meta-analysis. *Br J Sports Med* 48:973-+.
- Westerberg H, Jacobaeus H, Hirvikoski T, Clevberger P, Ostensson J, Bartfai A, Forssberg H, Klingberg T (2007) Computerized working memory training - a method of cognitive rehabilitation after stroke. *Brain Inj* 21:21-29.